

Interdisciplinarity of bioaerosol research/Bioaerosolu pētījumu starpdisciplināritāte



33rd International Scientific
Conference of the
University of Latvia 2025

Contribution ID: 1

Type: not specified

Long-Term Pollen Season Trends Of *Fraxinus*, *Quercus* And *Ambrosia artemisiifolia* As Indicators Of Anthropogenic Climate Change Impact

Thursday, 30 January 2025 14:30 (20 minutes)

Background

Climate change significantly affects plant life cycles, including flowering, particularly in anemophilous species, which can release high concentrations of allergenic pollen into the air. Changes in meteorological variables and air pollution can lead to earlier onset, prolonged duration, and increased intensity of pollen seasons, with adverse effects on individuals suffering from allergic diseases such as hay fever, rhinitis, or asthma. This study aimed to investigate long-term trends in pollen season characteristics for woody genera *Fraxinus* and *Quercus*, and herbaceous species *Ambrosia artemisiifolia*, and to explore their correlations with climatic and air pollution variables.

Methods

The research was conducted at two urban sites in Slovakia, Bratislava and Banská Bystrica (hereafter B. Bystrica), over the past two decades. Pollen was captured using Hirst-type traps placed at rooftop levels, and samples were analysed following standard aerobiological methods [1]. Daily mean pollen concentrations (pollen grains/m³) were calculated, and the main pollen season (MPS) was defined as the period between the first and last day when daily pollen concentrations reached 10 pollen grains/m³. Characteristics of the MPS were calculated, including start date, end date, duration, peak value, seasonal pollen integral, number of high days (above symptom-causing thresholds: 50 pollen/m³ for *Fraxinus* and *Quercus*, 20 pollen/m³ for *Ambrosia*), and the date of the first high day. Meteorological variables (mean, minimum, and maximum daily temperatures, precipitation, and relative humidity) and air pollutants (PM10, SO₂, NO₂, O₃, CO) were also analysed. To identify trends in these characteristics and variables, the Mann–Kendall trend test was employed. For variables with significant trends, the Theil–Sen estimator was applied to calculate the slope of the trend. Spearman’s correlation analysis was used to assess relationships between pollen season characteristics and significant trends in meteorological and air pollution variables.

Results

The study identified rising trends in temperature, precipitation, and air humidity, alongside mostly declining trends in air pollutants, except for increasing CO levels in B. Bystrica. For the woody taxa, pollen seasons demonstrated earlier onset, prolonged duration, and increased intensity, with *Quercus* showing more pronounced changes. For *Ambrosia*, trends included earlier start dates and extended durations at both locations and more high days and later end dates in B. Bystrica. Rising temperatures during the pre-season correlated with an earlier onset of the pollen season for trees, specifically significant for *Fraxinus* in Bratislava and *Quercus* in B. Bystrica. For *Ambrosia*, the significant delay in the end date of the pollen season correlated with increasing temperatures during its blooming period, particularly in B. Bystrica. Air pollutants, such as SO₂ and NO₂, negatively correlated with pollen intensity, while CO showed a positive correlation.

Conclusion

This study highlights the significant impact of climate change on the pollen seasons of allergenic plant taxa, emphasising the need for continued monitoring and prediction to mitigate health impacts on allergy sufferers. Understanding these trends contributes to a deeper comprehension of climate change effects on plant life cycles and aids in predicting future pollen seasons.

Acknowledgements

The authors acknowledge both the Division of Meteorology and Climatology (Faculty of Mathematics, Physics and Informatics of Comenius University in Bratislava) for providing meteorological data and the Slovak Hydrometeorological Institute for providing air pollution data used in this study.

Funding

This study was supported by Grant Agency VEGA (Bratislava), Grant No. 1/0180/22.

References

1. Galán, C., Cariñanos, P., Teno, A., & Domínguez, E. (2007). Spanish Aerobiology Network (REA): Management and Quality Manual. Servicio de Publicaciones, Universidad de Córdoba.

Please, submit you abstract

Primary authors: ŠČEVKOVÁ, Jana (Faculty of Natural Sciences, Department of Botany, Comenius University in Bratislava); ŠTEFÁNIKOVÁ, Natália (Faculty of Natural Sciences, Department of Botany, Comenius University in Bratislava); DUŠIČKA, Jozef (Faculty of Natural Sciences, Department of Botany, Comenius University in Bratislava); LAFFÉRSOVÁ, Janka (Department of Environmental Biology, Public Health Office, Banská Bystrica); ZAHRADNÍKOVÁ, Eva (Faculty of Natural Sciences, Department of Botany, Comenius University in Bratislava)

Presenter: ŠTEFÁNIKOVÁ, Natália (Faculty of Natural Sciences, Department of Botany, Comenius University in Bratislava)